

Measures for Reduction of Radionuclide Contamination of Agricultural Produce


April 2017

Ministry of Agriculture, Forestry and Fisheries


Table of contents

1 Control of radioactive materials in foods	2 – 4
2 Measures to reduce radionuclides migration	5 – 16
3 Results of inspection	17 – 19

1 Control of radioactive materials in foods



1 Control of radioactive materials in foods



1 Control of radioactive materials in foods

(Reference) Maximum levels for radioactive Cs in foods

- The maximum levels have been set in accordance with the table below

Food groups	Maximum level (Bq/kg)
Drinking water	10
Milk	50
General foods	100
Infant foods	50

- ※ Assuming that the intervention exemption level of 1 mSv/year, the same value as for Codex guideline levels.


4

2 Measures to reduce radionuclides migration

Pathways of radionuclide contamination of crops

Direct contamination by radionuclide fallout

Uptake of radionuclide from soil



5

2 Measures to reduce radionuclides migration

Measures for feed (Provisional tolerance values for radioactive Cs)

Feed for:	Set on 14 Apr. 2011	Revised on 1 Aug. 2011	Revised on 1 Apr. 2012
Cattle	300 Bq/kg ^{*1}	300 Bq/kg ^{*2}	100 Bq/kg ^{*2}
Pigs	-	300 Bq/kg ^{*1}	80 Bq/kg ^{*2}
Chickens	-	300 Bq/kg ^{*1}	160 Bq/kg ^{*2}
Cultured fish	-	100 Bq/kg ^{*3}	40 Bq/kg ^{*3}

* Provisional tolerance values for feed set on a basis of feed consumption and provisional transfer coefficients by:

*1 Referring the IAEA documents.

*2 Using the results of the transfer studies on dairy cattle, pigs and hens conducted by MAFF in Japan after the accident.

*3 Using the result of the cultured fish transfer study conducted by MAFF in Japan after the accident and previous studies.

6

2 Measures to reduce radionuclides migration

Feed management in accordance with tolerance values

1 Thorough enforcement of appropriate feed management by pastures and others under the provisional tolerance values.



2 Decontamination by deep plowing and others can lead to pasture production under provisionally tolerated level.



7

2 Measures to reduce radionuclides migration

Measures for materials used for the cultivation of edible fungi (Reference values for radioactive Cs)

Material	Set on 6 Oct. 2011	Revised on 1 Apr. 2012
Wood logs (for Shiitake)	150 Bq/kg	50 Bq/kg
Cultivation media	150 Bq/kg	200 Bq/kg

8

2 Measures to reduce radionuclides migration

Measures for the cultivation of edible fungi

- ❑ Introduction of safe production materials, and reduction of pollution due to radionuclides
- ❑ Information related to collection of wild edible fungi and wild edible plants

Efforts

1. Ensuring safe wood logs (purchase support for wood logs, supply and demand matching for wood logs)
2. Decontamination of wood logs, introduction of simple greenhouses, etc
3. Dissemination and guidance for cultivation management in accordance with guidelines
4. Dissemination of cultivation technology that reduces pollution from radionuclides
5. Transmission of information via websites and pamphlets, patrol guidance



9

2 Measures to reduce radionuclides migration

Measures for materials used as fertilizers, soil conditioners, and nursery soils (for radioactive Cs)

Material	Value	Note
Sludge for manure	200 Bq/kg	Standard value set on 24 Jun. 2011
Fertilizers (including those from leaves), soil conditioners, nursery soils, etc.	400 Bq/kg	Provisional tolerance value set on 1 Aug. 2011

10

2 Measures to reduce radionuclides migration

Measures to reduce radionuclides of fruit trees

The levels of radioactive Cs deposited onto above-ground parts of fruit trees have been reduced by cleaning the surface of bark with high-pressure water.

High-pressure washing of peach trees



Scraping bark for pear





11

2 Measures to reduce radionuclides migration

Measures to reduce radionuclides of tea trees

To prevent the transfer of radioactive Cs from leaves and branches to new leaves, leaves and branches were plucked or pruned further than in usual practice.




2 Measures to reduce radionuclides migration

Absorption control through potassic fertilization to rice

- ❑ Soil in paddy fields where rice with high-level radioactive Cs was produced tend to contain low-level potassium
- ❑ Having similar chemical characteristics to Cs, potassium in soil can suppress absorption of Cs by root uptake


[If potassium concentration in soil is suitable]



2 Measures to reduce radionuclides migration

(Reference) Application of potassium

(Test results in farmlands with over 500 Bq/kg)



14

2 Measures to reduce radionuclides migration

Decontamination of farmland (Removal to topsoil)

Remove the surface soil to remove the nuclides in soil surface



Results of removal of topsoil (2011, Iitatemura)

Removal of topsoil

Before: 10,370 Bq/kg

After: 2,599 Bq/kg (75% reduction)

Ambient dose level (Surface)

Before: 7.1 μ Sv/hr

After: 3.4 μ Sv/hr (52% reduction)


After harvest: 1.9 μ Sv/hr

15

2 Measures to reduce radionuclides migration

Decontamination of farmland (Deep plowing)

Deep plowing to replace top soil with subsoil to keep the most of fallen radionuclides deeper than the range of plant root



**Deep plowing
(30 cm)**

16

3 Results of Inspection

□ Number of samples

	Year (FY)				
	2011 ²⁾	2012	2013	2014	2015
Test samples Total ¹⁾	93,121	212,927	248,111	233,608	260,538
Available for cultivation/feed management ³⁾	83,520	190,431	222,690	207,511	237,845
Difficult for cultivation/feed management ⁴⁾	6,489	20,482	23,565	24,642	21,236
Others (Processed food, etc.)	3,112	2,014	1,856	1,455	1,457

1: Prefectural product pre-shipment inspection samples in 17 prefectures (Aomori, Iwate, Akita, Miyagi, Yamagata, Fukushima, Ibaraki, Tochigi, Gunma, Chiba, Saitama, Tokyo, Kanagawa, Niigata, Yamanashi, Nagano, Shizuoka)

2: Including March 2011

3: Vegetables, Tubers, Fruits, Seeds, Rice, Grains, Legumes, Cereals, Meat, Eggs, Fresh Milk, Tea (ready for consumption), Edible Fungi (cultivated), Wild Plants for Food (cultivated)

4: Wild edible fungi, Wild plants for Food (wild), Game meat, Fishery Products, Honey

17

3 Results of Inspection

□ Radioactive Cs concentration in foods

(**Available** for cultivation/feed management)

Radioactive Cs concentration (Bq/kg)	2011	2012	2013	2014	2015
Less than 25	79,073 (96.2%)	184,431 (98.5%)	217,770 (98.9%)	204,970 (99.8%)	235,881 (99.8%)
25 – 50	1,289 (1.6%)	1,242 (0.7%)	1,328 (0.6%)	380 (0.2%)	325 (0.1%)
50 – 75	696 (0.8%)	631 (0.3%)	509 (0.2%)	37 (0.02%)	34 (0.01%)
75 – 100	339 (0.4%)	500 (0.3%)	446 (0.2%)	9 (0.004%)	12 (0.01%)
More than 100	830 (1.0%)	399 (0.2%)	87 (0.04%)	5 (0.002%)	5 (0.002%)

1: Upper row: detected samples, lower row: detected samples rate for inspected samples

2: Excluding fresh milk and tea, for which maximum levels differ from general foodstuffs

18

3 Results of Inspection

□ Radioactive Cs concentration in foods

(**Difficult** for cultivation/feed management)

Radioactive Cs concentration (Bq/kg)	2011	2012	2013	2014	2015
Less than 25	3,565 (55%)	14,805 (72%)	20,332 (86%)	22,358 (91%)	20,080 (95%)
25 - 50	781 (12%)	2,237 (11%)	1,321 (5.6%)	1,065 (4.3%)	552 (2.6%)
50 - 75	473 (7.3%)	983 (4.8%)	617 (2.6%)	424 (1.7%)	224 (1.1%)
75 - 100	327 (5.0%)	635 (3.1%)	393 (1.7%)	253 (1.0%)	121 (0.6%)
More than 100	1,343 (21%)	1,822 (8.9%)	902 (3.8%)	542 (2.2%)	259 (1.2%)

Upper row: detected samples, lower row: detected samples rate for inspected samples

19