

# Measures for Reduction of Radionuclide Contamination of Agricultural Produce

June 2021

Ministry of Agriculture, Forestry and Fisheries

## Table of contents

1 Control of radioactive materials in foods	2 – 4
2 Measures to reduce radionuclides migration	5 – 16
3 Test Results	17 – 19

# 1 Control of radioactive materials in foods

## Nuclear Emergency Response Headquarters

- Restrict food shipments / Set and remove intake limits

Instruction ↓

### Related municipalities

- Develop and implement inspection plans for food
- Implementation of shipment and intake restrictions for foodstuffs

Claims / Support  
←  
→  
Reports

## Ministry of Health, Labor, and Welfare

### Set the maximum levels for radioactive materials in food

- Disclose test results

↑ Support

Cooperation ↔

Inquiries ↓ Findings ↑

## Ministry of Agriculture, Forestry, and Fisheries

- Regulate materials used for production of foods and feeds
- Support inspection plans and advise in technical inspection
- Advise in technical provisions for reduction of radionuclide contamination at production sites

## Food Safety Commission

- Evaluate health impact from radioactive materials in food

## Nuclear Regulation Authority

- Radiation Council

2

# 1 Control of radioactive materials in foods

Restriction on planting of crops,  
In particular, rice

Measures for reduction of  
radionuclide contamination

- Regulation of materials used for production of foods and feeds
- Decontamination of crops
- Decontamination of soils in farmland

Analysis of  
contamination factors

Sampling and testing

If exceed

Restriction of distribution

3

# 1 Control of radioactive materials in foods

(Reference) Maximum levels for radioactive Cs in foods

- The maximum levels have been set in accordance with the table below

Food groups	Maximum level (Bq/kg)
Drinking water	10
Milk	50
General foods	100
Infant foods	50

- ※ Assuming that the intervention exemption level of 1 mSv/year, the same value as for Codex guideline levels.



4

## 2 Measures to reduce radionuclides migration

### Pathways of radionuclide contamination of crops

Direct contamination by radionuclide fallout

Uptake of radionuclide from soil



Radionuclides attached to trees are transferred to fruits or shoot

5

## 2 Measures to reduce radionuclides migration

Measures for feed (Provisional tolerance values for radioactive Cs)

Feed for:	Set on 14 Apr. 2011	Revised on 1 Aug. 2011	Revised on 1 Apr. 2012
Cattle	300 Bq/kg* <sup>1</sup>	300 Bq/kg* <sup>2</sup>	100 Bq/kg* <sup>2</sup>
Pigs	-	300 Bq/kg* <sup>1</sup>	80 Bq/kg* <sup>2</sup>
Chickens	-	300 Bq/kg* <sup>1</sup>	160 Bq/kg* <sup>2</sup>
Cultured fish	-	100 Bq/kg* <sup>3</sup>	40 Bq/kg* <sup>3</sup>

\* Provisional tolerance values for feed set on a basis of feed consumption and provisional transfer coefficients by:

\*<sup>1</sup> Referring the IAEA documents.

\*<sup>2</sup> Using the results of the transfer studies on dairy cattle, pigs and hens conducted by MAFF in Japan after the accident.

\*<sup>3</sup> Using the result of the cultured fish transfer study conducted by MAFF in Japan after the accident and previous studies.

6

## 2 Measures to reduce radionuclides migration

Feed management in accordance with tolerance values

1 Thorough enforcement of appropriate feed management by pastures and others under the provisional tolerance values.



2 Decontamination by deep plowing and others can lead to pasture production under provisionally tolerated level.



7

## 2 Measures to reduce radionuclides migration

Measures for materials used for the cultivation of edible fungi (Reference values for radioactive Cs)

Material	Set on 6 Oct. 2011	Revised on 1 Apr. 2012
Wood logs (for Shiitake)	150 Bq/kg	50 Bq/kg
Cultivation media	150 Bq/kg	200 Bq/kg

8

## 2 Measures to reduce radionuclides migration

Measures for the cultivation of edible fungi

- ❑ Introduction of safe production materials, and reduction of pollution due to radionuclides
- ❑ Information related to collection of wild edible fungi and wild edible plants

### Efforts

1. Securing safe wood logs (wood logs procurement support and supply-demand matching for wood logs)
2. Support for decontamination of wood logs and introduction of pollution control facilities including simple greenhouses
3. Establishment and dissemination of wood log fungi cultivation management guidelines for radionuclides reduction
4. Dissemination of cultivation technology that reduces pollution from radionuclides
5. Transmission of information via websites and pamphlets, on-site traveling guidance



9



## 2 Measures to reduce radionuclides migration

Measures for materials used as fertilizers, soil conditioners, and nursery soils (for radioactive Cs)

Material	Value	Note
Sludge for manure	200 Bq/kg	Standard value set on 24 Jun. 2011
Fertilizers (including those from leaves), soil conditioners, nursery soils, etc.	400 Bq/kg	Provisional tolerance value set on 1 Aug. 2011

10

## 2 Measures to reduce radionuclides migration

### Measures to reduce radionuclides of fruit trees

The levels of radioactive Cs deposited onto above-ground parts of fruit trees have been reduced by cleaning the surface of bark with high-pressure water.

High-pressure washing of peach trees



Scraping bark for pear




11

## 2 Measures to reduce radionuclides migration

### Measures to reduce radionuclides of tea trees

To prevent the transfer of radioactive Cs from leaves and branches to new leaves, leaves and branches were plucked or pruned further than in usual practice.



Before pruning



After pruning




12

## 2 Measures to reduce radionuclides migration

### Absorption control through potassic fertilization to rice

- ❑ Soil in paddy fields where rice with high-level radioactive Cs was produced tend to contain low-level potassium
- ❑ Having similar chemical characteristics to Cs, potassium in soil can suppress absorption of Cs by root uptake

[If potassium concentration in soil is suitable]




13

## 2 Measures to reduce radionuclides migration

### (Reference) Application of potassium

(Test results in farmlands with over 500 Bq/kg)



14

## 2 Measures to reduce radionuclides migration

### Decontamination of farmland (Removal to topsoil)

Remove the surface soil to remove the nuclides in soil surface



### Results of removal of topsoil (2011, litatemura)

#### Removal of topsoil

Before: 10,370 Bq/kg

After: 2,599 Bq/kg (**75% reduction**)

#### Ambient dose level (Surface)

Before: 7.1  $\mu$ Sv/hr

After: 3.4  $\mu$ Sv/hr (**52% reduction**)

After harvest: 1.9  $\mu$ Sv/hr


15



## 2 Measures to reduce radionuclides migration

### Decontamination of farmland (Deep plowing)

Deep plowing to replace top soil with subsoil to keep the most of fallen radionuclides deeper than the range of plant root



**Deep plowing  
(30 cm)**

16

## 3 Test Results

### Number of samples

	Year (FY)									
	2011 <sup>2)</sup>	2012	2013	2014	2015	2016	2017	2018	2019	2020
Test samples Total <sup>1)</sup>	93,288	212,922	248,273	233,738	260,794	249,833	245,086	240,002	229,333	44,258
Available for cultivation/feed management <sup>3)</sup>	83,676	190,376	222,725	207,484	237,741	225,662	223,239	219,977	212,332	28,354
Difficult for cultivation/feed management <sup>4)</sup>	6,491	20,530	23,707	24,794	21,596	22,812	20,833	19,040	16,080	15,115
Others (Processed food, etc.)	3,121	2,016	1,841	1,460	1,457	1,359	1,014	985	921	789

1: Prefectural product pre-shipment test samples in 17 prefectures (Aomori, Iwate, Akita, Miyagi, Yamagata, Fukushima, Ibaraki, Tochigi, Gunma, Chiba, Saitama, Tokyo, Kanagawa, Niigata, Yamanashi, Nagano, Shizuoka)

2: Including March 2011

3: Vegetables, Tubers, Fruits, Seeds, Rice, Grains, Legumes, Cereals, Meat, Eggs, Fresh Milk, Tea(ready for consumption), Edible Fungi (cultivated), Wild Plants for Food (cultivated)

4: Wild edible fungi, Wild plants for Food (wild), Game meat, Fishery Products, Honey

17

## 3 Test Results

### Radioactive Cs concentration in foods

(**Available** for cultivation/feed management )

Vegetables, Tubers, Fruits, Seeds, Rice, Grains, Legumes, Cereals, Meat, Eggs, Edible Fungi (cultivated), Wild Plants for Food (cultivated)

Radioactive Cs concentration (Bq/kg)	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Less than 25	79,229 (96.17%)	184,396 (98.53%)	217,785 (98.92%)	204,941 (99.79%)	235,781 (99.84%)	223,779 (99.84%)	222,109 (99.90%)	219,084 (99.90%)	211,635 (99.93%)	27,945 (99.54%)
25–50	1,293 (1.57%)	1,237 (0.66%)	1,340 (0.61%)	382 (0.19%)	321 (0.14%)	332 (0.15%)	209 (0.09%)	194 (0.09%)	141 (0.07%)	116 (0.41%)
50–100	1036 (1.26%)	1123 (0.60%)	960 (0.44%)	45 (0.02%)	46 (0.02%)	31 (0.01%)	17 (0.01%)	30 (0.01%)	16 (0.01%)	14 (0.05%)
More than 100	825 (1.00%)	392 (0.21%)	87 (0.04%)	5 (0.002%)	5 (0.002%)	0 (0%)	2 (0.001%)	0 (0%)	2 (0.001%)	0 (0%)

1: Upper row: detected samples, lower row: detected samples rate for inspected samples

2: Excluding fresh milk and tea, for which maximum levels differ from general foodstuffs.

18

## 3 Test Results

### Radioactive Cs concentration in foods

(**Difficult** for cultivation/feed management )

Wild edible fungi, Wild plants for Food (wild), Game meat, Fishery Products, Honey

Radioactive Cs concentration (Bq/kg)	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Less than 25	3,567 (54.95%)	14,835 (72.26%)	20,455 (86.28%)	22,506 (90.77%)	20,412 (94.52%)	21,092 (92.46%)	19,855 (95.31%)	17,696 (92.94%)	15,207 (94.57%)	14,234 (94.17%)
25–50	781 (12.0%)	2,241 (10.92%)	1,336 (5.64%)	1,068 (4.31%)	573 (2.65%)	845 (3.70%)	534 (2.56%)	700 (3.68%)	541 (3.36%)	627 (4.15%)
50–100	800 (12.3%)	1,626 (7.92%)	1,014 (4.28%)	678 (2.73%)	352 (1.63%)	425 (1.86%)	259 (1.24%)	350 (1.84%)	175 (1.09%)	151 (1.00%)
More than 100	1,343 (20.69%)	1,828 (8.90%)	902 (3.80%)	542 (2.19%)	259 (1.20%)	450 (1.97%)	185 (0.89%)	294 (1.54%)	157 (0.98%)	103 (0.68%)

Upper row: detected samples, lower row: detected samples rate for inspected samples

19