

豚熱感染はイノシシの 行動様式に变化をもたらすのか

小寺祐二・宇都宮大学

CSF**未感染地域**

社会構造 行動様式


耳票型GPS




サーキットデザイン社：GLT-04


Table 3. Number of wild boars captured in Yamizo area during 2011 - 2012 and 2014 - 2016.
Normal figures are the number of individuals, and those in parentheses show the percentages each year.

Age (years)	Total	Year										Chi-square test
		2011		2012		2014		2015		2016		
3.0 over	9	0	(0.0)	2	(2.2)	2	(2.4)	2	(2.3)	3	(3.5)	$df = 24$ $P = 0.0048$
2.5	46	3	(6.3)	10	(10.9)	8	(9.8)	9	(10.3)	16	(18.6)	
2.0	62	6	(12.5)	12	(13.0)	11	(13.4)	11	(12.6)	22	(25.6)	
1.5	52	4	(8.3)	7	(7.6)	11	(13.4)	19	(21.8)	11	(12.8)	
1.0	102	11	(22.9)	22	(23.9)	26	(31.7)	19	(21.8)	24	(27.9)	
0.5	64	11	(22.9)	21	(22.8)	11	(13.4)	13	(14.9)	8	(9.3)	
0.0	60	13	(27.1)	18	(19.6)	13	(15.9)	14	(16.1)	2	(2.3)	
Total	395	48	(100.0)	92	(100.0)	82	(100.0)	87	(100.0)	86	(100.0)	












Image © 2025 Airbus
Data SIO, NOAA, U.S. Navy, NGA, GEBCO

Google Earth

画像取得日: 2024/11/13 36° 31'50.47" N 140° 13'49.82" E 標高 164 m 高度 100.65 km



Image © 2025 Airbus

Google Earth

画像取得日: 2023/5/17 36° 31'50.46" N 140° 13'49.82" E 標高 156 m 高度 2074 km