

SN: 11040001
玄米 RICE, HUSKED

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
37,250	< 0.01	1.2


カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	3,113	8.4	3,113	8.4
0.01 ≤ X ≤ 0.10	27,839	74.7	30,952	83.1
0.10 < X ≤ 0.20	5,057	13.6	36,009	96.7
0.20 < X ≤ 0.30	946	2.5	36,955	99.2
0.30 < X ≤ 0.40	201	0.5	37,156	99.7
0.40 < X ≤ 0.50	55	0.1	37,211	99.9
0.5 < X	39	0.1	37,250	100.0

注: 土壌試料は分析していない

b) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.2 mg/kg	3.3%
0.3 mg/kg	0.8%
0.4 mg/kg ^注	0.3%
0.5 mg/kg	0.1%

注: 下線の値は検討されている基準値原案。



SN: 11040002, 11040003, 11040004
その他穀類 OTHER CEREALS (inc. 大麦 BARLEY, 裸麦 RYE and そば BUCKWHEAT)

a) 収穫前調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
66	< 0.01	0.07


カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	11	16.7	11	16.7
0.01 ≤ X ≤ 0.10	55	83.3	66	100.0
0.10 < X ≤ 0.20	0			
0.20 < X ≤ 0.30	0			
0.30 < X ≤ 0.40	0			
0.40 < X ≤ 0.50	0			
0.5 < X	0			

注: 大麦、裸麦、そばの分析結果については、u-検定により母集団が同一であると判断されたため、度数分布表を統合している。

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
66	< 0.05	0.77

カドミウム濃度 (mg/kg)	試料数
X < 0.05	3
0.05 ≤ X ≤ 0.10	8
0.10 < X ≤ 0.20	20
0.20 < X ≤ 0.30	25
0.30 < X ≤ 0.40	8
0.40 < X ≤ 0.50	1
0.5 < X	1



SN: 11040005
小麦 WHEAT

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
382	< 0.01	0.47

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	5	1.3	5	1.3
0.01 ≤ X ≤ 0.10	292	76.4	297	77.7
0.10 < X ≤ 0.20	73	19.1	370	96.9
0.20 < X ≤ 0.30	9	2.4	379	99.2
0.30 < X ≤ 0.40	2	0.5	381	99.7
0.40 < X ≤ 0.50	1	0.3	382	100.0
0.5 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
227	< 0.05	0.8

カドミウム濃度 (mg/kg)	試料数
X < 0.05	27
0.05 ≤ X ≤ 0.10	34
0.10 < X ≤ 0.20	109
0.20 < X ≤ 0.30	36
0.30 < X ≤ 0.40	8
0.40 < X ≤ 0.50	8
0.5 < X	5

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
<u>0.2 mg/kg</u> 注	3.1%
0.3 mg/kg	0.8%
0.4 mg/kg	0.3%

注: 下線の値は検討されている基準値原案。



SN: 11040006
ゆりねLILY, BULB

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
23	0.03	0.17

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	0	0.0	0	0.0
0.01 ≤ X ≤ 0.05	7	30.4	7	30.4
0.05 < X ≤ 0.10	10	43.5	17	73.9
0.10 < X ≤ 0.15	4	17.4	21	91.3
0.15 < X ≤ 0.20	2	8.7	23	100.0
0.20 < X ≤ 0.25	0			
0.25 < X ≤ 0.30	0			
0.30 < X ≤ 0.35	0			
0.35 < X ≤ 0.40	0			
0.40 < X ≤ 0.45	0			
0.45 < X ≤ 0.5	0			
0.5 < X	0			

注: 土壌試料は分析していない。

SN: 11040007
にんにく GARLIC

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
95	< 0.01	0.20


カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	4	4.2	4	4.2
0.01 ≤ X ≤ 0.05	63	66.3	67	70.5
0.05 < X ≤ 0.10	20	21.1	87	91.6
0.10 < X ≤ 0.15	6	6.3	93	97.9
0.15 < X ≤ 0.20	2	2.1	95	100.0
0.20 < X ≤ 0.25	0			
0.25 < X ≤ 0.30	0			
0.30 < X ≤ 0.35	0			
0.35 < X ≤ 0.40	0			
0.40 < X ≤ 0.45	0			
0.45 < X ≤ 0.5	0			
0.5 < X	0			

注: 土壌試料は分析していない。

b) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.05 mg/kg ^注	29.5%
0.10 mg/kg	8.4%
0.20 mg/kg	0%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
105	< 0.01	0.07

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	50	47.6	50	47.6
0.01 ≤ X ≤ 0.02	34	32.4	84	80.0
0.02 < X ≤ 0.03	11	10.5	95	90.5
0.03 < X ≤ 0.04	7	6.7	102	97.1
0.04 < X ≤ 0.05	2	1.9	104	99.0
0.05 < X ≤ 0.06	0	0.0	104	99.0
0.06 < X ≤ 0.07	1	1.0	105	100.0
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
9	0.10	0.41

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	1
0.10 < X ≤ 0.20	6
0.20 < X ≤ 0.30	0
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	1
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値 ML values	超過率 Rate (%)
0.05 mg/kg 注	1.0%
0.10 mg/kg	0%
0.20 mg/kg	0%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
128	< 0.01	0.16

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	75	58.6	75	58.6
0.01 ≤ X ≤ 0.02	37	28.9	112	87.5
0.02 < X ≤ 0.03	5	3.9	117	91.4
0.03 < X ≤ 0.04	4	3.1	121	94.5
0.04 < X ≤ 0.05	2	1.6	123	96.1
0.05 < X ≤ 0.06	0	0.0	123	96.1
0.06 < X ≤ 0.07	1	0.8	124	96.9
0.07 < X ≤ 0.08	1	0.8	125	97.7
0.08 < X ≤ 0.09	2	1.6	127	99.2
0.09 < X ≤ 0.10	0	0.0	127	99.2
0.10 < X	1	0.8	128	100.0

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
66	< 0.05	0.57

カドミウム濃度 (mg/kg)	試料数
X < 0.05	6
0.05 ≤ X ≤ 0.10	17
0.10 < X ≤ 0.20	27
0.20 < X ≤ 0.30	13
0.30 < X ≤ 0.40	2
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	1
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.05 mg/kg 注	3.9%
0.10 mg/kg	0.8%
0.20 mg/kg	0.0%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
117	< 0.01	0.011

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	110	94.0	110	94.0
0.01 ≤ X ≤ 0.02	7	6.0	117	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
63	< 0.05	0.36

カドミウム濃度 (mg/kg)	試料数
X < 0.05	3
0.05 ≤ X ≤ 0.10	10
0.10 < X ≤ 0.20	29
0.20 < X ≤ 0.30	18
0.30 < X ≤ 0.40	3
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.05 mg/kg	0%
0.20 mg/kg ^注	0%

注: 下線の値は検討されている基準値原案。



SN: 11040011
ブロッコリー BROCCOLI

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
32	< 0.01	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	22	68.8	22	68.8
0.01 ≤ X ≤ 0.02	8	25.0	30	93.8
0.02 < X ≤ 0.03	2	6.3	32	100.0
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
16	0.1	0.35

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	2
0.10 < X ≤ 0.20	7
0.20 < X ≤ 0.30	6
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040012
カリフラワー CAULIFLOWER

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
20	< 0.01	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	18	90.0	18	90.0
0.01 ≤ X ≤ 0.02	1	5.0	19	95.0
0.02 < X ≤ 0.03	0	0.0	19	95.0
0.03 < X ≤ 0.04	1	5.0	20	100.0
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

SN: 11040013

メロン MELONS, EXCEPT WATERMELON

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
24	< 0.01	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	18	75.0	18	75.0
0.01 ≤ X ≤ 0.02	6	25.0	24	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
8	0.08	0.35

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	1
0.10 < X ≤ 0.20	3
0.20 < X ≤ 0.30	3
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
84	< 0.01	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	82	97.6	82	97.6
0.01 ≤ X ≤ 0.02	2	2.4	84	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
29	0.07	0.48

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	5
0.10 < X ≤ 0.20	12
0.20 < X ≤ 0.30	6
0.30 < X ≤ 0.40	3
0.40 < X ≤ 0.50	3
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.02 mg/kg	0%
0.05 mg/kg 注	0%

注: 下線の値は検討されている基準値原案。



SN: 11040015

すいか WATERMELON

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
40	< 0.01	< 0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	40	100.0	40	100.0
0.01 ≤ X ≤ 0.02	0			
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
20	< 0.05	0.45

カドミウム濃度 (mg/kg)	試料数
X < 0.05	2
0.05 ≤ X ≤ 0.10	3
0.10 < X ≤ 0.20	8
0.20 < X ≤ 0.30	3
0.30 < X ≤ 0.40	3
0.40 < X ≤ 0.50	1
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040016

かぼちゃ WINTER SQUASH

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
24	< 0.01	0.011

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	22	91.7	22	91.7
0.01 ≤ X ≤ 0.02	2	8.3	24	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
4	0.09	0.27

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	2
0.10 < X ≤ 0.20	1
0.20 < X ≤ 0.30	1
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

a) 収穫前及び収穫後調査


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
594	< 0.01	0.66

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	5	0.8	5	0.8
0.01 ≤ X ≤ 0.10	284	47.8	289	48.7
0.10 < X ≤ 0.20	202	34.0	491	82.7
0.20 < X ≤ 0.30	66	11.1	557	93.8
0.30 < X ≤ 0.40	26	4.4	583	98.1
0.40 < X ≤ 0.50	7	1.2	590	99.3
0.50 < X ≤ 0.60	2	0.3	592	99.7
0.60 < X ≤ 0.70	2	0.3	594	100.0
0.70 < X ≤ 0.80	0			
0.8 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
427	< 0.05	0.60

カドミウム濃度 (mg/kg)	試料数
X < 0.05	18
0.05 ≤ X ≤ 0.10	64
0.10 < X ≤ 0.20	217
0.20 < X ≤ 0.30	88
0.30 < X ≤ 0.40	28
0.40 < X ≤ 0.50	7
0.50 < X ≤ 0.60	5
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0



SN: 11040018

小豆 ADZUKI BEAN, DRY

a) 収穫前調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
14	< 0.01	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	13	92.9	13	92.9
0.01 ≤ X ≤ 0.10	1	7.1	14	100.0
0.10 < X ≤ 0.20	0			
0.20 < X ≤ 0.30	0			
0.30 < X ≤ 0.40	0			
0.40 < X ≤ 0.50	0			
0.5 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
14	0.07	0.26

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	3
0.10 < X ≤ 0.20	10
0.20 < X ≤ 0.30	1
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040019

みつば MITSUBA

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
18	< 0.01	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	15	83.3	15	83.3
0.01 ≤ X ≤ 0.02	3	16.7	18	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
42	< 0.01	0.12

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	8	19.0	8	19.0
0.01 ≤ X ≤ 0.02	13	31.0	21	50.0
0.02 < X ≤ 0.04	8	19.0	29	69.0
0.04 < X ≤ 0.06	3	7.1	32	76.2
0.06 < X ≤ 0.08	4	9.5	36	85.7
0.08 < X ≤ 0.10	3	7.1	39	92.9
0.10 < X ≤ 0.12	3	7.1	42	100.0
0.12 < X ≤ 0.14	0			
0.14 < X ≤ 0.16	0			
0.16 < X ≤ 0.18	0			
0.18 < X ≤ 0.20	0			
0.20 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
12	0.11	0.4

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	0
0.10 < X ≤ 0.20	5
0.20 < X ≤ 0.30	2
0.30 < X ≤ 0.40	5
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
51	< 0.01	0.09

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	11	21.6	11	21.6
0.01 ≤ X ≤ 0.02	18	35.3	29	56.9
0.02 < X ≤ 0.03	8	15.7	37	72.5
0.03 < X ≤ 0.04	7	13.7	44	86.3
0.04 < X ≤ 0.05	3	5.9	47	92.2
0.05 < X ≤ 0.06	1	2.0	48	94.1
0.06 < X ≤ 0.07	1	2.0	49	96.1
0.07 < X ≤ 0.08	0	0.0	49	96.1
0.08 < X ≤ 0.09	2	3.9	51	100.0
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: こまつなとみずなの分析結果については、u-検定による母集団が同一であると判定されたため、度数分布表を統合している。

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
20	0.08	0.35

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	7
0.10 < X ≤ 0.20	7
0.20 < X ≤ 0.30	3
0.30 < X ≤ 0.40	3
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040022

チンゲンサイ PAK-CHOI

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
23	< 0.01	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	9	40.9	9	40.9
0.01 ≤ X ≤ 0.02	10	40.9	19	81.8
0.02 < X ≤ 0.03	3	13.6	22	95.5
0.03 < X ≤ 0.04	1	4.5	23	100.0
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
110	< 0.01	0.056

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	58	52.7	58	52.7
0.01 ≤ X ≤ 0.02	47	42.7	105	95.5
0.02 < X ≤ 0.03	2	1.8	107	97.3
0.03 < X ≤ 0.04	2	1.8	109	99.1
0.04 < X ≤ 0.05	0	0.0	109	99.1
0.05 < X ≤ 0.06	1	0.9	110	100.0
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
39	< 0.05	0.34

カドミウム濃度 (mg/kg)	試料数
X < 0.05	5
0.05 ≤ X ≤ 0.10	6
0.10 < X ≤ 0.20	15
0.20 < X ≤ 0.30	11
0.30 < X ≤ 0.40	2
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.05 mg/kg	0.9%
0.10 mg/kg	0%
0.20 mg/kg _注	0%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
90	< 0.01	0.08

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	22	24.4	22	24.4
0.01 ≤ X ≤ 0.02	49	54.4	71	78.9
0.02 < X ≤ 0.03	10	11.1	81	90.0
0.03 < X ≤ 0.04	6	6.7	87	96.7
0.04 < X ≤ 0.05	1	1.1	88	97.8
0.05 < X ≤ 0.06	1	1.1	89	98.9
0.06 < X ≤ 0.07	0	0.0	89	98.9
0.07 < X ≤ 0.08	1	1.1	90	100.0
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
29	< 0.05	0.42

カドミウム濃度 (mg/kg)	試料数
X < 0.05	1
0.05 ≤ X ≤ 0.10	1
0.10 < X ≤ 0.20	12
0.20 < X ≤ 0.30	12
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	2
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.05 mg/kg	2.3%
0.10 mg/kg	0%
<u>0.20 mg/kg</u> 注	0%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
435	< 0.01	0.49

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	14	3.2	14	3.2
0.01 ≤ X ≤ 0.10	370	85.1	384	88.3
0.10 < X ≤ 0.20	38	8.7	422	97.0
0.20 < X ≤ 0.30	9	2.1	431	99.1
0.30 < X ≤ 0.40	2	0.5	433	99.5
0.40 < X ≤ 0.50	2	0.5	435	100.0
0.50 < X ≤ 0.60	0			
0.60 < X ≤ 0.70	0			
0.70 < X ≤ 0.80	0			
0.80 < X ≤ 0.90	0			
0.90 < X ≤ 1.0	0			
1.0 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
73	< 0.05	0.37

カドミウム濃度 (mg/kg)	試料数
X < 0.05	3
0.05 ≤ X ≤ 0.10	14
0.10 < X ≤ 0.20	40
0.20 < X ≤ 0.30	13
0.30 < X ≤ 0.40	3
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.2 mg/kg 注	3.0%
0.3 mg/kg	0.9%
0.4 mg/kg	0.5%
0.5 mg/kg	0%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
400	< 0.01	0.17

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	167	41.8	167	41.8
0.01 ≤ X ≤ 0.02	137	34.3	304	76.0
0.02 < X ≤ 0.03	41	10.3	345	86.3
0.03 < X ≤ 0.04	22	5.5	367	91.8
0.04 < X ≤ 0.05	10	2.5	377	94.3
0.05 < X ≤ 0.06	7	1.8	384	96.0
0.06 < X ≤ 0.07	5	1.3	389	97.3
0.07 < X ≤ 0.08	3	0.8	392	98.0
0.08 < X ≤ 0.09	3	0.8	395	98.8
0.09 < X ≤ 0.10	2	0.5	397	99.3
0.10 < X	3	0.8	400	100.0

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
66	< 0.05	0.5

カドミウム濃度 (mg/kg)	試料数
X < 0.05	1
0.05 ≤ X ≤ 0.10	4
0.10 < X ≤ 0.20	27
0.20 < X ≤ 0.30	18
0.30 < X ≤ 0.40	12
0.40 < X ≤ 0.50	4
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.05 mg/kg 注	5.8%
0.1 mg/kg	0.7%
0.2 mg/kg	0%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
165	< 0.01	0.22

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	3	1.8	3	1.8
0.01 ≤ X ≤ 0.02	35	21.2	38	23.0
0.02 < X ≤ 0.03	43	26.1	81	49.1
0.03 < X ≤ 0.04	26	15.8	107	64.8
0.04 < X ≤ 0.05	21	12.7	128	77.6
0.05 < X ≤ 0.06	13	7.9	141	85.5
0.06 < X ≤ 0.07	5	3.0	146	88.5
0.07 < X ≤ 0.08	7	4.2	153	92.7
0.08 < X ≤ 0.09	1	0.6	154	93.3
0.09 < X ≤ 0.10	6	3.6	160	97.0
0.10 < X	5	3.0	165	100.0

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
10	< 0.05	0.18

カドミウム濃度 (mg/kg)	試料数
X < 0.05	3
0.05 ≤ X ≤ 0.10	4
0.10 < X ≤ 0.20	3
0.20 < X ≤ 0.30	0
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.05 mg/kg 注	22.4%
0.1 mg/kg	3.0%
0.2 mg/kg	0.6%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
130	< 0.01	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	44	33.8	44	33.8
0.01 ≤ X ≤ 0.02	61	46.9	105	80.8
0.02 < X ≤ 0.03	20	15.4	125	96.2
0.03 < X ≤ 0.04	5	3.8	130	100.0
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
38	< 0.05	0.33

カドミウム濃度 (mg/kg)	試料数
X < 0.05	4
0.05 ≤ X ≤ 0.10	2
0.10 < X ≤ 0.20	22
0.20 < X ≤ 0.30	8
0.30 < X ≤ 0.40	2
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.02 mg/kg	19.2%
0.05 mg/kg 注	0%

注: 下線の値は検討されている基準値原案。



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
137	< 0.01	0.05

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	92	67.2	92	67.2
0.01 ≤ X ≤ 0.02	37	27.0	129	94.2
0.02 < X ≤ 0.03	6	4.4	135	98.5
0.03 < X ≤ 0.04	0	0.0	135	98.5
0.04 < X ≤ 0.05	2	1.5	137	100.0
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			


b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
55	< 0.05	0.39

カドミウム濃度 (mg/kg)	試料数
X < 0.05	2
0.05 ≤ X ≤ 0.10	12
0.10 < X ≤ 0.20	23
0.20 < X ≤ 0.30	11
0.30 < X ≤ 0.40	7
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.02 mg/kg	5.4%
0.05 mg/kg	0%



a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
32	< 0.01	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	27	84.4	27	84.4
0.01 ≤ X ≤ 0.02	4	12.5	31	96.9
0.02 < X ≤ 0.03	1	3.1	32	100.0
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
11	< 0.05	0.35

カドミウム濃度 (mg/kg)	試料数
X < 0.05	1
0.05 ≤ X ≤ 0.10	2
0.10 < X ≤ 0.20	3
0.20 < X ≤ 0.30	2
0.30 < X ≤ 0.40	3
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040032
未成熟空豆 BROAD BEEN (GREEN PODS AND IMMATURE SEEDS)

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
19	< 0.01	0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	18	94.7	18	94.7
0.01 ≤ X ≤ 0.02	1	5.3	1	5.3
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

SN: 11040033
さやいんげん COMMON BEEN (PODS AND IMMATURE SEEDS)

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
21	C < 0.01	C < 0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	21	100.0	21	100.0
0.01 ≤ X ≤ 0.02	0			
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

SN: 11040034
さやえんどう GARDEN PEA (YOUNG PODS)

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
22	< 0.01	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	果積試料数	割合 (%)
X < 0.01	21	95.5	21	95.5
0.01 ≤ X ≤ 0.02	1	4.5	22	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
5	0.08	0.39

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	1
0.10 < X ≤ 0.20	3
0.20 < X ≤ 0.30	0
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040035
グリーンピース GARDEN PEA, SHELLED

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
14	< 0.01	0.05

カドミウム濃度 (mg/kg)	試料数	割合 (%)	果積試料数	割合 (%)
X < 0.01	8	57.1	8	57.1
0.01 ≤ X ≤ 0.02	5	35.7	13	92.9
0.02 < X ≤ 0.03	0	0.0	13	92.9
0.03 < X ≤ 0.04	0	0.0	13	92.9
0.04 < X ≤ 0.05	1	7.1	14	100.0
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
24	< 0.01	0.05

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	7	29.2	7	29.2
0.01 ≤ X ≤ 0.02	16	66.7	23	95.8
0.02 < X ≤ 0.03	1	4.2	24	100.0
0.03 < X ≤ 0.04	0	0.0	24	100.0
0.04 < X ≤ 0.05	1	4.2	25	104.2
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
3	0.21	0.30

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	0
0.10 < X ≤ 0.20	0
0.20 < X ≤ 0.30	3
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
20	< 0.01	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	16	80.0	16	80.0
0.01 ≤ X ≤ 0.02	4	20.0	20	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
35	< 0.01	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	22	62.9	22	62.9
0.01 ≤ X ≤ 0.02	13	37.1	35	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
3	0.07	0.21

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	1
0.10 < X ≤ 0.20	0
0.20 < X ≤ 0.30	2
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

a) 収穫前調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
6	< 0.01	0.066

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	1	16.7	1	16.7
0.01 ≤ X ≤ 0.02	2	33.3	3	50.0
0.02 < X ≤ 0.03	1	16.7	4	66.7
0.03 < X ≤ 0.04	1	16.7	5	83.3
0.04 < X ≤ 0.05	0	0.0	5	83.3
0.05 < X ≤ 0.06	0	0.0	5	83.3
0.06 < X ≤ 0.07	1	16.7	6	100.0
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注: 土壌試料は分析していない

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
6	0.09	0.35

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	1
0.10 < X ≤ 0.20	2
0.20 < X ≤ 0.30	2
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
302	< 0.01	0.33

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	40	13.2	40	13.2
0.01 ≤ X ≤ 0.05	188	62.3	228	75.5
0.05 < X ≤ 0.10	44	14.6	272	90.1
0.10 < X ≤ 0.15	17	5.6	289	95.7
0.15 < X ≤ 0.20	9	3.0	298	98.7
0.20 < X ≤ 0.25	1	0.3	299	99.0
0.25 < X ≤ 0.30	2	0.7	301	99.7
0.30 < X ≤ 0.35	1	0.3	302	100.0
0.35 < X ≤ 0.40	0			
0.40 < X ≤ 0.45	0			
0.45 < X ≤ 0.50	0			
0.5 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
36	< 0.05	0.62

カドミウム濃度 (mg/kg)	試料数
X < 0.05	1
0.05 ≤ X ≤ 0.10	4
0.10 < X ≤ 0.20	19
0.20 < X ≤ 0.30	6
0.30 < X ≤ 0.40	4
0.40 < X ≤ 0.50	1
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	1
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.1 mg/kg ^注	9.9%
0.2 mg/kg	1.3%
0.3 mg/kg	0.3%
0.5 mg/kg	0.0%

注: 下線の値は検討されている基準値原案。



SN: 11040041

甘藷 SWEET POTATO

a) 収穫前調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
37	< 0.01	0.014

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	31	83.8	31	83.8
0.01 ≤ X ≤ 0.02	6	16.2	37	100.0
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 収穫後調査(皮をむいたもの)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
40	< 0.01	0.050

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	32	80.0	32	80.0
0.01 ≤ X ≤ 0.02	5	12.5	37	92.5
0.02 < X ≤ 0.03	0	0.0	37	92.5
0.03 < X ≤ 0.04	2	5.0	39	97.5
0.04 < X ≤ 0.05	1	2.5	40	100.0
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

c) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
37	< 0.05	0.27

カドミウム濃度 (mg/kg)	試料数
X < 0.05	14
0.05 ≤ X ≤ 0.10	4
0.10 < X ≤ 0.20	15
0.20 < X ≤ 0.30	4
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)		最大値 (mg/kg)	
125	< 0.01		0.23	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	9	7.2	9	7.2
0.01 ≤ X ≤ 0.05	84	67.2	93	74.4
0.05 < X ≤ 0.10	25	20.0	118	94.4
0.10 < X ≤ 0.15	6	4.8	124	99.2
0.15 < X ≤ 0.20	0	0.0	124	99.2
0.20 < X ≤ 0.25	1	0.8	125	100.0
0.25 < X ≤ 0.30	0			
0.30 < X ≤ 0.35	0			
0.35 < X ≤ 0.40	0			
0.40 < X ≤ 0.45	0			
0.45 < X ≤ 0.50	0			
0.5 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
8	< 0.05	0.23

カドミウム濃度 (mg/kg)	試料数
X < 0.05	1
0.05 ≤ X ≤ 0.10	2
0.10 < X ≤ 0.20	4
0.20 < X ≤ 0.30	1
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.1 mg/kg ^注	5.6%
0.2 mg/kg	0.8%
0.3 mg/kg	0%

注: 下線の値は検討されている基準値原案。



a) Field samples

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
35	< 0.01	0.07


カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	11	31.4	11	31.4
0.01 ≤ X ≤ 0.05	22	62.9	33	94.3
0.05 < X ≤ 0.10	2	5.7	35	100.0
0.10 < X ≤ 0.15	0			
0.15 < X ≤ 0.20	0			
0.20 < X ≤ 0.25	0			
0.25 < X ≤ 0.30	0			
0.30 < X ≤ 0.35	0			
0.35 < X ≤ 0.40	0			
0.40 < X ≤ 0.45	0			
0.45 < X ≤ 0.50	0			
0.5 < X	0			

b) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
134	< 0.01	0.16

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	16	11.9	16	11.9
0.01 ≤ X ≤ 0.05	105	78.4	121	90.3
0.05 < X ≤ 0.10	11	8.2	132	98.5
0.10 < X ≤ 0.15	1	0.7	133	99.3
0.15 < X ≤ 0.20	1	0.7	134	100.0
0.20 < X ≤ 0.25	0			
0.25 < X ≤ 0.30	0			
0.30 < X ≤ 0.35	0			
0.35 < X ≤ 0.40	0			
0.40 < X ≤ 0.45	0			
0.45 < X ≤ 0.50	0			
0.5 < X	0			

注: 収穫前調査と収穫後調査の結果は、z-検定により母集団が異なると判断されたため、度数分布表を分けて作成している。



にんじん(つづき)

c) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
35	< 0.05	0.44

カドミウム濃度 (mg/kg)	試料数
X < 0.05	3
0.05 ≤ X ≤ 0.10	5
0.10 < X ≤ 0.20	21
0.20 < X ≤ 0.30	4
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	1
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

d) 想定基準値に対する超過率

想定基準値	超過率 (%)
<u>0.1 mg/kg</u> 注	1.5%
0.2 mg/kg	0%

注: 下線の値は検討されている基準値原案。

SN: 11040044

馬鈴薯 POTATO

a) 収穫前調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
29	< 0.01	0.058

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	2	6.9	2	6.9
0.01 ≤ X ≤ 0.02	10	34.5	12	41.4
0.02 < X ≤ 0.03	5	17.2	17	58.6
0.03 < X ≤ 0.04	9	31.0	26	89.7
0.04 < X ≤ 0.05	2	6.9	28	96.6
0.05 < X ≤ 0.06	1	3.4	29	100.0
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 収穫後調査(皮をむいたもの)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
40	< 0.01	0.040

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	6	15.0	6	15.0
0.01 ≤ X ≤ 0.02	20	50.0	26	65.0
0.02 < X ≤ 0.03	7	17.5	33	82.5
0.03 < X ≤ 0.04	7	17.5	40	100.0
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

c) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
29	< 0.05	0.17

カドミウム濃度 (mg/kg)	試料数
X < 0.05	8
0.05 ≤ X ≤ 0.10	4
0.10 < X ≤ 0.20	17
0.20 < X ≤ 0.30	0
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040045

だいこん RADISH, JAPANESE

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
107	< 0.01	0.050

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	81	75.7	81	75.7
0.01 ≤ X ≤ 0.02	25	23.4	106	99.1
0.02 < X ≤ 0.03	0	0.0	106	99.1
0.03 < X ≤ 0.04	0	0.0	106	99.1
0.04 < X ≤ 0.05	1	0.9	107	100.0
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
54	< 0.05	0.27

カドミウム濃度 (mg/kg)	試料数
X < 0.05	18
0.05 ≤ X ≤ 0.10	16
0.10 < X ≤ 0.20	17
0.20 < X ≤ 0.30	3
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
0.05 mg/kg	0%
0.1 mg/kg 注	0%

注: 下線の値は検討されている基準値原案。



a) 収穫前調査


全試料数	最小値 (mg/kg)	最大値 (mg/kg)
17	< 0.01	0.10

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	6	35.3	6	35.3
0.01 ≤ X ≤ 0.05	8	47.1	14	82.4
0.05 < X ≤ 0.10	3	17.6	17	100.0
0.10 < X ≤ 0.15	0			
0.15 < X ≤ 0.20	0			
0.20 < X ≤ 0.25	0			
0.25 < X ≤ 0.30	0			
0.30 < X ≤ 0.35	0			
0.35 < X ≤ 0.40	0			
0.40 < X ≤ 0.45	0			
0.45 < X ≤ 0.50	0			
0.5 < X	0			

b) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
53	< 0.01	0.18

Concentration (mg/kg)	No. of samples	Ratio (%)	Cumulative No. of samples	Ratio (%)
X < 0.01	22	41.5	22	41.5
0.01 ≤ X ≤ 0.05	24	45.3	46	86.8
0.05 < X ≤ 0.10	6	11.3	52	98.1
0.10 < X ≤ 0.15	0	0.0	52	98.1
0.15 < X ≤ 0.20	1	1.9	53	100.0
0.20 < X ≤ 0.25	0			
0.25 < X ≤ 0.30	0			
0.30 < X ≤ 0.35	0			
0.35 < X ≤ 0.40	0			
0.40 < X ≤ 0.45	0			
0.45 < X ≤ 0.50	0			
0.5 < X	0			



やまいも(つづき)

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
17	< 0.05	0.27

カドミウム濃度 (mg/kg)	試料数
X < 0.05	3
0.05 ≤ X ≤ 0.10	6
0.10 < X ≤ 0.20	7
0.20 < X ≤ 0.30	1
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

c) 想定基準値に対する超過率

想定基準値	超過率 (%)
<u>0.1 mg/kg</u> 注	1.9%
0.2 mg/kg	0%

注: 下線の値は検討されている基準値原案。

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
3	0.012	0.031

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	0	0.0	0	0.0
0.01 ≤ X ≤ 0.02	2	66.7	2	66.7
0.02 < X ≤ 0.03	0	0.0	2	66.7
0.03 < X ≤ 0.04	1	33.3	3	100.0
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
3	0.26	0.27

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	0
0.10 < X ≤ 0.20	0
0.20 < X ≤ 0.30	3
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

a) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
30	< 0.01	0.07

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	10	33.3	10	33.3
0.01 ≤ X ≤ 0.02	11	36.7	21	70.0
0.02 < X ≤ 0.03	5	16.7	26	86.7
0.03 < X ≤ 0.04	1	3.3	27	90.0
0.04 < X ≤ 0.05	1	3.3	28	93.3
0.05 < X ≤ 0.06	1	3.3	29	96.7
0.06 < X ≤ 0.07	1	3.3	30	100.0
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

注:土壌試料は調査していない。

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
41	< 0.01	0.08

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	18	43.9	18	43.9
0.01 ≤ X ≤ 0.02	16	39.0	34	82.9
0.02 < X ≤ 0.03	4	9.8	38	92.7
0.03 < X ≤ 0.04	1	2.4	39	95.1
0.04 < X ≤ 0.05	0	0.0	39	95.1
0.05 < X ≤ 0.06	1	2.4	40	97.6
0.06 < X ≤ 0.07	0	0.0	40	97.6
0.07 < X ≤ 0.08	1	2.4	41	100.0
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
11	0.07	0.33

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	1
0.10 < X ≤ 0.20	4
0.20 < X ≤ 0.30	5
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040050

セレリー CELERY

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
26	< 0.01	0.08

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	4	15.4	4	15.4
0.01 ≤ X ≤ 0.02	8	30.8	12	46.2
0.02 < X ≤ 0.03	5	19.2	17	65.4
0.03 < X ≤ 0.04	4	15.4	21	80.8
0.04 < X ≤ 0.05	3	11.5	24	92.3
0.05 < X ≤ 0.06	0	0.0	24	92.3
0.06 < X ≤ 0.07	1	3.8	25	96.2
0.07 < X ≤ 0.08	1	3.8	26	100.0
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
7	0.14	0.33

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	0
0.10 < X ≤ 0.20	4
0.20 < X ≤ 0.30	2
0.30 < X ≤ 0.40	1
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040052
いちご STRAWBERRY

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
50	< 0.01	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	37	74.0	37	74.0
0.01 ≤ X ≤ 0.02	11	22.0	48	96.0
0.02 < X ≤ 0.03	1	2.0	49	98.0
0.03 < X ≤ 0.04	1	2.0	50	100.0
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
18	0.12	0.49

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	0
0.10 < X ≤ 0.20	6
0.20 < X ≤ 0.30	7
0.30 < X ≤ 0.40	4
0.40 < X ≤ 0.50	1
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040056
なし NASHI PEAR

a) 収穫前調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
43	< 0.01	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	32	74.4	32	74.4
0.01 ≤ X ≤ 0.02	9	20.9	41	95.3
0.02 < X ≤ 0.03	2	4.7	43	100.0
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
20	< 0.01	< 0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	20	100.0	20	100.0
0.01 ≤ X ≤ 0.02	0			
0.02 < X ≤ 0.03	0			
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

c) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
44	0.07	0.62

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	6
0.10 < X ≤ 0.20	15
0.20 < X ≤ 0.30	15
0.30 < X ≤ 0.40	6
0.40 < X ≤ 0.50	1
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	1
0.70 < X ≤ 0.80	0

以下の試料については、分析結果がいずれも定量下限値を下回った。(0.01 mg/kg)

ぶどう 30試料
りんご 39試料
桃 16試料
さくらんぼ 1試料
柿 16試料
キウイフルーツ 3試料
柑橘類 79試料

SN: 11040060

にら CHIVES, CHINESE

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
23	< 0.01	0.16

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	9	39.1	9	39.1
0.01 ≤ X ≤ 0.02	7	30.4	16	69.6
0.02 < X ≤ 0.04	5	21.7	21	91.3
0.04 < X ≤ 0.06	1	4.3	22	95.7
0.06 < X ≤ 0.08	0	0.0	22	95.7
0.08 < X ≤ 0.10	0	0.0	22	95.7
0.10 < X ≤ 0.12	0	0.0	22	95.7
0.12 < X ≤ 0.14	0	0.0	22	95.7
0.14 < X ≤ 0.16	1	4.3	23	100.0
0.16 < X ≤ 0.18	0			
0.18 < X ≤ 0.20	0			
0.20 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
4	< 0.05	0.16

カドミウム濃度 (mg/kg)	試料数
X < 0.05	1
0.05 ≤ X ≤ 0.10	0
0.10 < X ≤ 0.20	3
0.20 < X ≤ 0.30	0
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040061

しょうが GINGER, ROOT

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
25	< 0.01	0.040

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	5	20.0	5	20.0
0.01 ≤ X ≤ 0.02	13	52.0	18	72.0
0.02 < X ≤ 0.03	4	16.0	22	88.0
0.03 < X ≤ 0.04	3	12.0	25	100.0
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
5	0.12	0.17

カドミウム濃度 (mg/kg)	試料数
X < 0.05	0
0.05 ≤ X ≤ 0.10	0
0.10 < X ≤ 0.20	5
0.20 < X ≤ 0.30	0
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

SN: 11040062

落花生 PEANUT

a) 収穫前及び収穫後調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
36	0.04	0.18

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	0	0.0	0	0.0
0.01 ≤ X ≤ 0.10	25	69.4	25	69.4
0.10 < X ≤ 0.20	11	30.6	36	100.0
0.20 < X ≤ 0.30	0			
0.30 < X ≤ 0.40	0			
0.40 < X ≤ 0.50	0			
0.50 < X	0			

SN: 11040063
栗 CHESTNUTS

a) 収穫前調査

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
4	< 0.01	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
X < 0.01	1	25.0	1	25.0
0.01 ≤ X ≤ 0.02	1	25.0	2	50.0
0.02 < X ≤ 0.03	2	50.0	4	100.0
0.03 < X ≤ 0.04	0			
0.04 < X ≤ 0.05	0			
0.05 < X ≤ 0.06	0			
0.06 < X ≤ 0.07	0			
0.07 < X ≤ 0.08	0			
0.08 < X ≤ 0.09	0			
0.09 < X ≤ 0.10	0			
0.10 < X	0			

b) 土壌試料

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
4	< 0.05	0.17

カドミウム濃度 (mg/kg)	試料数
X < 0.05	1
0.05 ≤ X ≤ 0.10	1
0.10 < X ≤ 0.20	2
0.20 < X ≤ 0.30	0
0.30 < X ≤ 0.40	0
0.40 < X ≤ 0.50	0
0.50 < X ≤ 0.60	0
0.60 < X ≤ 0.70	0
0.70 < X ≤ 0.80	0
0.8 < X	0

肉類

a) 牛肉 (MEAT OF CATTLE)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
116	< 0.01	0.05

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01 ^{注1}	110	94.8	110	94.8
C<0.02 ^{注2}	110	94.8	110	94.8
0.02 C<0.03	1	0.9	111	95.7
0.03 C<0.04	2	1.7	113	97.4
0.04 C<0.05	1	0.9	114	98.3
0.05 C	2	1.7	116	100.0

b) 豚肉 (MEAT OF PIG)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
121	< 0.01	0.07

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01 ^{注1}	116	95.9	116	95.9
C<0.02 ^{注2}	116	95.9	116	95.9
0.02 C<0.03	2	1.7	118	97.5
0.03 C<0.04	0	0.0	118	97.5
0.04 C<0.05	1	0.8	119	98.3
0.05 C	2	1.7	121	100.0

注1: 試料の濃度が検出限界よりも小さいことを示す。

注2: 試料の濃度が、添加回収試験で用いた標準溶液の最も低い濃度よりも小さいことを示す。

c) 鶏肉 (MEAT OF POULTRY)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
26	< 0.01	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01 ^{注1}	23	88.5	23	88.5
C<0.02 ^{注2}	23	88.5	23	88.5
0.02 C<0.03	1	0.9	24	92.3
0.03 C<0.04	2	1.7	26	100.0
0.04 C<0.05	0			
0.05 C	0			

d) 馬肉 (MEAT OF HORSE)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)
7	< 0.01	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01 ^{注1}	6	85.7	6	85.7
C<0.02 ^{注2}	6	85.7	6	85.7
0.02 C<0.03	0	0.0	6	85.7
0.03 C<0.04	1	0.9	7	100.0
0.04 C<0.05	0			
0.05 C	0			

1. 魚類
アユ (Smelt (Ayu))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
24	C<0.01	0.05	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	11	45.8	11	45.8
0.01 C 0.1	13	54.2	24	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

イシガレイ (Stone flounder)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
6	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	6	100.0	6	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

イシモチ (White croaker)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 < C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ウグイ (White mullet)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
12	C<0.01	0.02	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	10	83.3	10	83.3
0.01 < C 0.1	2	16.7	12	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ウナギ (Japanese eel)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	15	100.0	15	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

カツオ (Skipjack tuna)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	C<0.01	0.04	0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	10	66.7	10	66.7
0.01 C 0.1	5	33.3	15	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

キハダ (Yellowfin tuna)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	C<0.01	0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	4	80.0	4	80.0
0.01 C 0.1	1	20.0	5	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

クロカジキ (Pacific blue marlin)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	0.01	0.07	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	5	100.0	5	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

クロダイ (Black seabream)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
3	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 < C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

クロマグロ (Bluefin tuna)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
5	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	5	100.0	5	100.0
0.01 < C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

コイ (Common carp)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
24	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	24	100.0	24	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

コイチ (White flower croaker)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
3	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

コノシロ(Threashfin shad)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 < C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

シロザケ(Chum salmon)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
18	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	18	100.0	18	100.0
0.01 < C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

スケソウダラ (Alaska pollack)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
15	C<0.01	0.02	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	13	86.7	13	86.7
0.01 C 0.1	2	13.3	15	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

スズキ (Japanese seabass)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
24	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	24	100.0	24	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

タチウオ (Largehead hairtail)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
3	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ニジマス (Rainbow trout)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
15	C<0.01	0.02	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	11	73.3	11	73.3
0.01 C 0.1	4	26.7	15	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

パシウカジキ (Pacific sailfish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	0.06	0.35	0.14

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 < C 0.1	3	60.0	3	60.0
0.1 < C 0.2	1	20.0	4	80.0
0.2 < C 0.3	0	0.0	4	80.0
0.3 < C 0.4	1	20.0	5	100.0
0.4 < C 0.5	0			
0.5 < C 0.6	0			
0.6 < C 0.7	0			
0.7 < C 0.8	0			
0.8 < C 0.9	0			
0.9 < C 1.0	0			
1.0 < C 1.1	0			
1.1 < C 1.2	0			
1.2 < C 1.3	0			
1.3 < C 1.4	0			
1.4 < C 1.5	0			
1.5 < C 1.6	0			
1.6 < C 1.7	0			
1.7 < C 1.8	0			
1.8 < C 1.9	0			
1.9 < C 2.0	0			
2.0 < C 2.1	0			
2.1 < C 2.2	0			
2.2 < C 2.3	0			
2.3 < C 2.4	0			
2.4 < C 2.5	0			
2.5 < C 2.6	0			
2.6 < C 2.7	0			
2.7 < C 2.8	0			
2.8 < C 2.9	0			
2.9 < C 3.0	0			
3.0 < C	0			

ハタハタ (Japanese sandfish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	0.02	0.02	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 < C 0.1	3	100.0	3	100.0
0.1 < C 0.2	0			
0.2 < C 0.3	0			
0.3 < C 0.4	0			
0.4 < C 0.5	0			
0.5 < C 0.6	0			
0.6 < C 0.7	0			
0.7 < C 0.8	0			
0.8 < C 0.9	0			
0.9 < C 1.0	0			
1.0 < C 1.1	0			
1.1 < C 1.2	0			
1.2 < C 1.3	0			
1.3 < C 1.4	0			
1.4 < C 1.5	0			
1.5 < C 1.6	0			
1.6 < C 1.7	0			
1.7 < C 1.8	0			
1.8 < C 1.9	0			
1.9 < C 2.0	0			
2.0 < C 2.1	0			
2.1 < C 2.2	0			
2.2 < C 2.3	0			
2.3 < C 2.4	0			
2.4 < C 2.5	0			
2.5 < C 2.6	0			
2.6 < C 2.7	0			
2.7 < C 2.8	0			
2.8 < C 2.9	0			
2.9 < C 3.0	0			
3.0 < C	0			

ヒラメ (Flatfish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
18	C<0.01	0.02	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	16	88.9	16	88.9
0.01 C 0.1	2	11.1	18	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ビンナガ (Albacore)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
5	C<0.01	0.02	0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	1	20.0	1	20.0
0.01 C 0.1	4	80.0	5	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

フナ (Crucian carp)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
6	C<0.01	0.03	0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	50.0	3	50.0
0.01 C 0.1	3	50.0	6	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ブリ (Yellowtail)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
18	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	18	100.0	18	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マアジ(Horse-mackerel)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	15	100.0	15	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マアナゴ(Conger eel)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
21	C<0.01	0.03	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	15	71.4	15	71.4
0.01 C 0.1	6	28.6	21	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マイワシ (Sardin)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
15	C<0.01	0.03	0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	11	73.3	11	73.3
0.01 C 0.1	4	26.7	15	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マコガレイ (flounder)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
15	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	15	100.0	15	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マサバ(Chub-mackerel)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
18	C<0.01	0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	17	94.4	17	94.4
0.01 C 0.1	1	5.6	18	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マダイ (Red sea-bream)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
18	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	18	100.0	18	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マハゼ (Spiny goby)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ミナミマグロ (Southern bluefin tuna)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	0.01	0.06	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	5	100.0	5	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

メカジキ (Swordfish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	0.02	0.04	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	5	100.0	5	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

メバチ (Bigeye tuna)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	C<0.01	0.02	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	1	20.0	1	20.0
0.01 C 0.1	4	80.0	5	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

メバル(Brown rockfish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
3	C<0.01	C<0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ヤツメウナギ (Japanese lamprey)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
3	0.03	0.14	0.10	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	1	33.3	1	33.3
0.1<C 0.2	2	66.7	3	100.0
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ヨシキリザメ (Great blue shark)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	C<0.01	0.05	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	1	20.0	1	20.0
0.01 C 0.1	4	80.0	5	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ワカサギ (Smelt)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
21	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	21	100.0	21	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

2. 甲殻類
ガザミ (Common shoe-crab)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
30	C<0.01	0.29	0.07

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	8	26.7	8	26.7
0.01 < C 0.1	15	50.0	23	76.7
0.1<C 0.2	5	16.7	28	93.3
0.2<C 0.3	2	6.7	30	100.0
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

クルマエビ (Kuruma-prawn)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
35	C<0.01	0.41	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	17	48.6	17	48.6
0.01 < C 0.1	15	42.9	32	91.4
0.1<C 0.2	1	2.9	33	94.3
0.2<C 0.3	1	2.9	34	97.1
0.3<C 0.4	0	0.0	34	97.1
0.4<C 0.5	1	2.9	35	100.0
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ケガニ (Korean crab (Muscle))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
30	0.02	0.17	0.08

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	24	80.0	24	80.0
0.1<C 0.2	6	20.0	30	100.0
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

スジエビ (Freshwater prawn)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
18	0.02	0.11	0.05

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	17	94.4	17	94.4
0.1<C 0.2	1	5.6	18	100.0
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

テナガエビ (Scampi)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 C 0.1	0			
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ベニズワイガニ (Red snow crab)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
30	0.04	0.48	0.16

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	11	36.7	11	36.7
0.1<C 0.2	9	30.0	20	66.7
0.2<C 0.3	8	26.7	28	93.3
0.3<C 0.4	1	3.3	29	96.7
0.4<C 0.5	1	3.3	30	100.0
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ホッコクアカエビ (Deepwater prawn)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
45	0.02	0.57	0.11

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	33	73.3	33	73.3
0.1<C 0.2	5	11.1	38	84.4
0.2<C 0.3	2	4.4	40	88.9
0.3<C 0.4	4	8.9	44	97.8
0.4<C 0.5	0	0.0	44	97.8
0.5<C 0.6	1	2.2	45	100.0
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

3. 軟体動物
アカガイ (Chest-shell)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	0.60	0.68	0.64

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	0	0.0	0	0.0
0.1<C 0.2	0	0.0	0	0.0
0.2<C 0.3	0	0.0	0	0.0
0.3<C 0.4	0	0.0	0	0.0
0.4<C 0.5	0	0.0	0	0.0
0.5<C 0.6	1	33.3	1	33.3
0.6<C 0.7	2	66.7	3	100.0
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

アサリ (Japanese short-neck)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
51	0.02	0.17	0.06

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	49	96.1	49	96.1
0.1<C 0.2	2	3.9	51	100.0
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

アワビ (Abalone(muscle))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	0.02	0.07	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	15	100.0	15	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

イイダコ (Ocellated octopus)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
3	0.04	0.05	0.04	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	3	100.0	3	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

コウイカ (Cuttlefish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
15	C<0.01	0.01	C<0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	1	6.7	1	6.7
0.01 C 0.1	14	93.3	15	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

サザエ (Horned turban(Muscle))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
15	C<0.01	0.10	0.05	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	2	13.3	2	13.3
0.01 C 0.1	13	86.7	15	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

シジミ (Japanese corbicula)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
64	0.03	0.77	0.37	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	7	10.9	7	10.9
0.1<C 0.2	11	17.2	18	28.1
0.2<C 0.3	7	10.9	25	39.1
0.3<C 0.4	21	32.8	46	71.9
0.4<C 0.5	0	0.0	46	71.9
0.5<C 0.6	0	0.0	46	71.9
0.6<C 0.7	9	14.1	55	85.9
0.7<C 0.8	9	14.1	64	100.0
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

スルメイカ (Pacific flying squid)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
56	0.03	1.30	0.29

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	18	32.1	18	32.1
0.1<C 0.2	11	19.6	29	51.8
0.2<C 0.3	9	16.1	38	67.9
0.3<C 0.4	6	10.7	44	78.6
0.4<C 0.5	4	7.1	48	85.7
0.5<C 0.6	1	1.8	49	87.5
0.6<C 0.7	1	1.8	50	89.3
0.7<C 0.8	1	1.8	51	91.1
0.8<C 0.9	0	0.0	51	91.1
0.9<C 1.0	2	3.6	53	94.6
1.0<C 1.1	2	3.6	55	98.2
1.1<C 1.2	0	0.0	55	98.2
1.2<C 1.3	1	1.8	56	100.0
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ハマグリ (Japanese hard clam)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
48	0.02	0.14	0.07

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	40	83.3	40	83.3
0.1<C 0.2	8	16.7	48	100.0
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ホタテ貝柱 (Common scallop (Muskelsehne))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
57	0.01	0.51	0.12

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	42	73.7	42	73.7
0.1<C 0.2	7	12.3	49	86.0
0.2<C 0.3	1	1.8	50	87.7
0.3<C 0.4	3	5.3	53	93.0
0.4<C 0.5	2	3.5	55	96.5
0.5<C 0.6	2	3.5	57	100.0
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マガキ (Giant pacific oyster)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
45	0.10	0.68	0.30

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	1	2.2	1	2.2
0.1<C 0.2	16	35.6	17	37.8
0.2<C 0.3	8	17.8	25	55.6
0.3<C 0.4	7	15.6	32	71.1
0.4<C 0.5	8	17.8	40	88.9
0.5<C 0.6	2	4.4	42	93.3
0.6<C 0.7	3	6.7	45	100.0
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

マダコ (Octopus)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
24	C<0.01	0.07	0.01	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	14	58.3	14	58.3
0.01 C 0.1	10	41.7	24	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

4 . 棘皮動物
ウニ (Sea urchin)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
45	0.02	0.34	0.17	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	8	17.8	8	17.8
0.1<C 0.2	24	53.3	32	71.1
0.2<C 0.3	10	22.2	42	93.3
0.3<C 0.4	3	6.7	45	100.0
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

ナマコ (Sea cucumber)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	C<0.01	0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	14	93.3	14	93.3
0.01 C 0.1	1	6.7	15	100.0
0.1<C 0.2	0			
0.2<C 0.3	0			
0.3<C 0.4	0			
0.4<C 0.5	0			
0.5<C 0.6	0			
0.6<C 0.7	0			
0.7<C 0.8	0			
0.8<C 0.9	0			
0.9<C 1.0	0			
1.0<C 1.1	0			
1.1<C 1.2	0			
1.2<C 1.3	0			
1.3<C 1.4	0			
1.4<C 1.5	0			
1.5<C 1.6	0			
1.6<C 1.7	0			
1.7<C 1.8	0			
1.8<C 1.9	0			
1.9<C 2.0	0			
2.0<C 2.1	0			
2.1<C 2.2	0			
2.2<C 2.3	0			
2.3<C 2.4	0			
2.4<C 2.5	0			
2.5<C 2.6	0			
2.6<C 2.7	0			
2.7<C 2.8	0			
2.8<C 2.9	0			
2.9<C 3.0	0			
3.0<C	0			

5. 魚介類内臓
アカニシ (内臓含む) (Thomas's rapa whelk)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	1.50	3.80	2.53

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	0	0.0	0	0.0
0.1<C 0.2	0	0.0	0	0.0
0.2<C 0.3	0	0.0	0	0.0
0.3<C 0.4	0	0.0	0	0.0
0.4<C 0.5	0	0.0	0	0.0
0.5<C 0.6	0	0.0	0	0.0
0.6<C 0.7	0	0.0	0	0.0
0.7<C 0.8	0	0.0	0	0.0
0.8<C 0.9	0	0.0	0	0.0
0.9<C 1.0	0	0.0	0	0.0
1.0<C 1.1	0	0.0	0	0.0
1.1<C 1.2	0	0.0	0	0.0
1.2<C 1.3	0	0.0	0	0.0
1.3<C 1.4	1	33.3	1	33.3
1.4<C 1.5	0	0.0	1	33.3
1.5<C 1.6	0	0.0	1	33.3
1.6<C 1.7	0	0.0	1	33.3
1.7<C 1.8	0	0.0	1	33.3
1.8<C 1.9	0	0.0	1	33.3
1.9<C 2.0	0	0.0	1	33.3
2.0<C 2.1	0	0.0	1	33.3
2.1<C 2.2	0	0.0	1	33.3
2.2<C 2.3	1	33.3	2	66.7
2.3<C 2.4	0	0.0	2	66.7
2.4<C 2.5	0	0.0	2	66.7
2.5<C 2.6	0	0.0	2	66.7
2.6<C 2.7	0	0.0	2	66.7
2.7<C 2.8	0	0.0	2	66.7
2.8<C 2.9	0	0.0	2	66.7
2.9<C 3.0	0	0.0	2	66.7
3.0<C	1	33.3	3	100.0

アワビ内臓 (Abalone (internal organ))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
15	2.20	5.60	3.11	
カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.5	0	0.0	0	0.0
0.5<C 1.0	0	0.0	0	0.0
1.0<C 1.5	0	0.0	0	0.0
1.5<C 2.0	0	0.0	0	0.0
2.0<C 2.5	1	6.7	1	6.7
2.5<C 3.0	8	53.3	9	60.0
3.0<C 3.5	4	26.7	13	86.7
3.5<C 4.0	1	6.7	14	93.3
4.0<C 4.5	0	0.0	14	93.3
4.5<C 5.0	0	0.0	14	93.3
5.0<C 5.5	0	0.0	14	93.3
5.5<C 6.0	1	6.7	15	100.0
6.0<C 6.5	0			
6.5<C 7.0	0			
7.0<C 7.5	0			
7.5<C 8.0	0			
8.0<C 8.5	0			
8.5<C 9.0	0			
9.0<C 9.5	0			
9.5<C 10	0			
10<C	0			

ガザミみそ (Common shoe-crab (internal organ))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)	
30	0.09	1.90	0.69	
カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.1	2	6.7	2	6.7
0.1<C 0.2	6	20.0	8	26.7
0.2<C 0.3	4	13.3	12	40.0
0.3<C 0.4	2	6.7	14	46.7
0.4<C 0.5	0	0.0	14	46.7
0.5<C 0.6	0	0.0	14	46.7
0.6<C 0.7	0	0.0	14	46.7
0.7<C 0.8	3	10.0	17	56.7
0.8<C 0.9	2	6.7	19	63.3
0.9<C 1.0	5	16.7	24	80.0
1.0<C 1.1	0	0.0	24	80.0
1.1<C 1.2	0	0.0	24	80.0
1.2<C 1.3	3	10.0	27	90.0
1.3<C 1.4	0	0.0	27	90.0
1.4<C 1.5	1	3.3	28	93.3
1.5<C 1.6	1	3.3	29	96.7
1.6<C 1.7	0	0.0	29	96.7
1.7<C 1.8	0	0.0	29	96.7
1.8<C 1.9	1	3.3	30	100.0
1.9<C 2.0	0			
2.0<C	0			

ケガニみそ (Korean crab (internal organ))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	0.79	3.50	1.99

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.5	0	0.0	0	0.0
0.5<C 1.0	1	6.7	1	6.7
1.0<C 1.5	4	26.7	5	33.3
1.5<C 2.0	2	13.3	7	46.7
2.0<C 2.5	5	33.3	12	80.0
2.5<C 3.0	2	13.3	14	93.3
3.0<C 3.5	1	6.7	15	100.0
3.5<C 4.0	0			
4.0<C 4.5	0			
4.5<C 5.0	0			
5.0<C 5.5	0			
5.5<C 6.0	0			
6.0<C 6.5	0			
6.5<C 7.0	0			
7.0<C 7.5	0			
7.5<C 8.0	0			
8.0<C 8.5	0			
8.5<C 9.0	0			
9.0<C 9.5	0			
9.5<C 10	0			
10<C	0			

サザエ内臓 (Horned turban (internal organ))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	1.20	9.50	4.65

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.5	0	0.0	0	0.0
0.5<C 1.0	0	0.0	0	0.0
1.0<C 1.5	5	33.3	5	33.3
1.5<C 2.0	0	0.0	5	33.3
2.0<C 2.5	0	0.0	5	33.3
2.5<C 3.0	1	6.7	6	40.0
3.0<C 3.5	0	0.0	6	40.0
3.5<C 4.0	1	6.7	7	46.7
4.0<C 4.5	1	6.7	8	53.3
4.5<C 5.0	1	6.7	9	60.0
5.0<C 5.5	0	0.0	9	60.0
5.5<C 6.0	1	6.7	10	66.7
6.0<C 6.5	0	0.0	10	66.7
6.5<C 7.0	1	6.7	11	73.3
7.0<C 7.5	0	0.0	11	73.3
7.5<C 8.0	1	6.7	12	80.0
8.0<C 8.5	1	6.7	13	86.7
8.5<C 9.0	0	0.0	13	86.7
9.0<C 9.5	2	13.3	15	100.0
9.5<C 10	0			
10<C	0			

スルメイカ肝臓 (Pacific flying squid (liver))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
41	6.60	96.00	33.90

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 10	8	19.5	8	19.5
10<C 20	9	22.0	17	41.5
20<C 30	6	14.6	23	56.1
30<C 40	5	12.2	28	68.3
40<C 50	4	9.8	32	78.0
50<C 60	1	2.4	33	80.5
60<C 70	3	7.3	36	87.8
70<C 80	2	4.9	38	92.7
80<C 90	2	4.9	40	97.6
90<C 100	1	2.4	41	100.0
100<C	0			

ベニズワイガニ内臓 (Red snow crab (internal organ))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	2.30	23.00	11.74

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 2.0	0	0.0	0	0.0
2.0<C 4.0	2	13.3	2	13.3
4.0<C 6.0	2	13.3	4	26.7
6.0<C 8.0	1	6.7	5	33.3
8.0<C 10	1	6.7	6	40.0
10 <C 12	1	6.7	7	46.7
12 <C 14	3	20.0	10	66.7
14 <C 16	0	0.0	10	66.7
16 <C 18	4	26.7	14	93.3
18 <C 20	0	0.0	14	93.3
20 <C 22	0	0.0	14	93.3
22 <C 24	1	6.7	15	100.0
24 <C 26	0			
26 <C 28	0			
28 <C 30	0			
30 < C	0			

ホタテウロ (Common scallop (mid-gut gland))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
72	1.30	16.00	5.80

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.5	0	0.0	0	0.0
0.5<C 1.0	0	0.0	0	0.0
1.0<C 1.5	4	5.6	4	5.6
1.5<C 2.0	8	11.1	12	16.7
2.0<C 2.5	1	1.4	13	18.1
2.5<C 3.0	5	6.9	18	25.0
3.0<C 3.5	1	1.4	19	26.4
3.5<C 4.0	1	1.4	20	27.8
4.0<C 4.5	4	5.6	24	33.3
4.5<C 5.0	6	8.3	30	41.7
5.0<C 5.5	6	8.3	36	50.0
5.5<C 6.0	6	8.3	42	58.3
6.0<C 6.5	1	1.4	43	59.7
6.5<C 7.0	4	5.6	47	65.3
7.0<C 7.5	7	9.7	54	75.0
7.5<C 8.0	6	8.3	60	83.3
8.0<C 8.5	2	2.8	62	86.1
8.5<C 9.0	1	1.4	63	87.5
9.0<C 9.5	2	2.8	65	90.3
9.5<C 10	3	4.2	68	94.4
10<C 10.5	0	0.0	68	94.4
10.5<C 11	0	0.0	68	94.4
11<C 11.5	0	0.0	68	94.4
11.5<C 12	1	1.4	69	95.8
12<C 12.5	0	0.0	69	95.8
12.5<C 13	0	0.0	69	95.8
13<C 13.5	0	0.0	69	95.8
13.5<C 14	1	1.4	70	97.2
14<C 14.5	0	0.0	70	97.2
14.5<C 15	1	1.4	71	98.6
15<C	1	1.4	72	100.0

ホタテ卵・生殖腺 (Common scallop (gonad))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
27	0.32	4.90	1.78

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 C 0.5	3	11.1	3	11.1
0.5<C 1.0	5	18.5	8	29.6
1.0<C 1.5	6	22.2	14	51.9
1.5<C 2.0	6	22.2	20	74.1
2.0<C 2.5	2	7.4	22	81.5
2.5<C 3.0	1	3.7	23	85.2
3.0<C 3.5	0	0.0	23	85.2
3.5<C 4.0	2	7.4	25	92.6
4.0<C 4.5	1	3.7	26	96.3
4.5<C 5.0	1	3.7	27	100.0
5.0<C 5.5	0			
5.5<C 6.0	0			
6.0<C 6.5	0			
6.5<C 7.0	0			
7.0<C 7.5	0			
7.5<C 8.0	0			
8.0<C 8.5	0			
8.5<C 9.0	0			
9.0<C 9.5	0			
9.5<C 10	0			
10<C	0			

6. 塩辛類
イカ塩辛

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg / kg)
30	0.09	9.90	2.63

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	累積割合 (%)
C < 0.05	0	0.0	0	0.0
0.05 < C 0.5	2	6.7	2	6.7
0.5<C 1.0	8	26.7	10	33.3
1.0<C 1.5	5	16.7	15	50.0
1.5<C 2.0	2	6.7	17	56.7
2.0<C 2.5	2	6.7	19	63.3
2.5<C 3.0	1	3.3	20	66.7
3.0<C 3.5	2	6.7	22	73.3
3.5<C 4.0	1	3.3	23	76.7
4.0<C 4.5	1	3.3	24	80.0
4.5<C 5.0	2	6.7	26	86.7
5.0<C 5.5	0	0.0	26	86.7
5.5<C 6.0	1	3.3	27	90.0
6.0<C 6.5	1	3.3	28	93.3
6.5<C 7.0	0	0.0	28	93.3
7.0<C 7.5	1	3.3	29	96.7
7.5<C 8.0	0	0.0	29	96.7
8.0<C 8.5	0	0.0	29	96.7
8.5<C 9.0	0	0.0	29	96.7
9.0<C 9.5	0	0.0	29	96.7
9.5<C 10.0	1	3.3	30	100.0

イカ塩辛(黒づくり)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg / kg)
5	1.40	6.50	3.26

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	累積割合 (%)
C < 0.05	0	0.0	0	0.0
0.05 < C 1.0	0	0.0	0	0.0
1.0<C 1.5	1	20.0	1	20.0
1.5<C 2.0	1	20.0	2	40.0
2.0<C 2.5	1	20.0	3	60.0
2.5<C 3.0	0	0.0	3	60.0
3.0<C 3.5	0	0.0	3	60.0
3.5<C 4.0	1	20.0	4	80.0
4.0<C 4.5	0	0.0	4	80.0
4.5<C 5.0	0	0.0	4	80.0
5.0<C 5.5	0	0.0	4	80.0
5.5<C 6.0	0	0.0	4	80.0
6.0<C 6.5	1	20.0	5	100.0

ウニ塩辛

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg / kg)	
6	C < 0.05	0.21	0.09	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	累積割合 (%)
C < 0.05	3	50.0	3	50.0
0.01 C 0.1	0	0.0	3	50.0
0.1<C 0.2	2	33.3	5	83.3
0.2<C 0.3	1	16.7	6	100.0

カツオ塩辛

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg / kg)	
10	0.17	1.10	0.64	

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	累積割合 (%)
C < 0.05	0	0.0	0	0.0
0.05 C 0.1	0	0.0	0	0.0
0.1<C 0.2	1	10.0	1	10.0
0.2<C 0.3	1	10.0	2	20.0
0.3<C 0.4	2	20.0	4	40.0
0.4<C 0.5	0	0.0	4	40.0
0.5<C 0.6	1	10.0	5	50.0
0.6<C 0.7	0	0.0	5	50.0
0.7<C 0.8	2	20.0	7	70.0
0.8<C 0.9	0	0.0	7	70.0
0.9<C 1.0	1	10.0	8	80.0
1.0<C 1.1	2	20.0	10	100.0